

NEW EVIDENCE FOR BLACK HOLE EVENT HORIZONS FROM CHANDRA

MICHAEL R. GARCIA,¹ JEFFREY E. MCCLINTOCK,¹ RAMESH NARAYAN,¹ PAUL CALLANAN,² & STEPHEN S. MURRAY¹

Submitted to ApJ Letters Dec 20, 2000

ABSTRACT

Previously we claimed that Black Hole X-ray Novae (BHXN) in quiescence are much less luminous than equivalent Neutron Star X-ray Novae (NSXN). This claim was based on the quiescent detection of a single short period BHXN (A0620–00, $P_{orb}=7.8$ hrs) and two longer period BHXN (GRO J1655–40, $P_{orb}=62.9$ hrs; V404 Cyg, $P_{orb}=155.3$ hrs), along with sensitive upper limits. We announce the detection of two more short period BHXN (GRO J0422+32, $P_{orb}=5.1$ hrs; GS 2000+25, $P_{orb}=8.3$ hrs), an upper limit for a third which is improved by two orders of magnitude (4U 1543–47, $P_{orb}=27.0$ hrs) and a new, much lower quiescent measurement of GRO J1655–40. Taken together, these new Chandra measurements confirm that the quiescent X-ray luminosities of BHXN are significantly lower than those of NSXN. We argue that this provides strong evidence for the existence of event horizons in BHXN.

Subject headings: black hole physics — X-ray: stars — binaries: close — stars: neutron — stars: individual (GRO J0422+32, A0620–00, GS 2000+25, 4U 1543–47, GRO J1655–40, and V404 Cyg)

1. INTRODUCTION

The very low, but non-zero, quiescent X-ray luminosity of the BHXN A0620–00 is difficult to understand in the context of standard viscous accretion disk theory (McClintock, Horne & Remillard 1995), given the continued mass transfer from the companion evidenced by an optically bright disk. The observations are explained by an advection-dominated accretion flow model, or ADAF (Narayan, McClintock & Yi 1996).

In an ADAF (see Narayan, Mahadevan & Quataert 1998 and Kato, Fukue & Mineshige 1998 for reviews), the energy released by accretion is stored as heat in a radiatively inefficient flow. Should the accreting object be a black hole this energy would be lost from view once it crosses the event horizon, but if the object has a solid surface the energy would be released upon impact with that surface and radiated to infinity. Thus, for the same mass accretion rate, a black hole would be significantly less luminous than a compact star with a surface (Narayan & Yi 1995). Such a comparison of otherwise similar systems is a promising method for proving the reality of event horizons (Narayan, Garcia & McClintock 1997a, hereafter NGM).

X-ray novae (XN) containing black hole (BH) and neutron star (NS) primaries are believed to be similar in many respects. Most importantly, the mass transfer rate from the secondary, measured in Eddington units of the primary, is believed to be comparable in BHXN and NSXN of similar orbital periods (Menou et al 1999, hereafter M99). NGM, Garcia et al. 1998 (hereafter G98) and M99 showed that quiescent BHXN are much less luminous in X-rays than quiescent NSXN of similar orbital periods, and argued that this provides direct evidence that BHXN are able to “hide” their accretion energy behind an event horizon. However, their claims were based on a very small sample in which only one detected BHXN (A0620–00, $P_{orb}=7.8$ hrs) had an orbital period similar to those of the comparison NSXN.

We report here new observations of four BHXN as part of the Chandra AO1 HRC Team GTO program and an additional two

BHXN as part of a complementary AO1 GO program.

2. OBSERVATIONS AND ANALYSIS

The basic parameters of the Chandra observations described here are presented in Table 1. Data were analyzed with a combination of the CXC CIAO V1.1 (CXC 2000a), HEASARC XSPEC V11.0 (Arnaud & Dorman 2000), and software written by Alexey Vikhlinin (Vikhlinin et al 1998). Detected fluxes and counting rates were measured over a 0.3–7.0 keV energy band for ACIS-S observations, and 0.5–7.0 keV for ACIS-I observations, in order to reduce the instrumental background. The observed on-axis PSF is such that 95% of the source flux is contained in the 1.5'' radius source extraction circle (van Speybroeck et al 1997; CXC 2000b). We observed occasional intervals of enhanced background during the observation of A0620–00 (see Plucinsky & Virani 2000), and therefore rejected 6.7 ks of this observation. The total accumulated background within the source extraction circle is < 1 count per source for all the observations.

We defer a full discussion of the Chandra spectra of quiescent BHXN to a second paper (McClintock et al 2001). Here we merely note that all the Chandra spectra are consistent with a single power-law with $\alpha \sim 2$ and absorption consistent with the optical extinction. We assume this in estimating the luminosities quoted here (unless explicitly stated otherwise).

GRO J0422+32.—The GRO J0422+32 region was observed with the Chandra Observatory (Weisskopf & O’Dell 1997) ACIS-I CCD array (Garmire et al 1992) for 18.8 ks on 2000 Oct 10. The CCD array recorded 16 photons at a position consistent with the optical position of GRO J0422+32, which constitutes a clear detection. Assuming a distance of 2.6 kpc (M99) and X-ray absorption corresponding to the optical extinction of $A_V = 1.2$ (Filippenko, Matheson, & Ho 1995; Predehl & Schmitt 1995), we derive an emitted luminosity of $7.6 \times 10^{30} \text{ erg s}^{-1}$ (0.5–10.0 keV). Luminosities, distances, primary masses, and $\log N_H$ for all sources are listed in Table 1.

A0620–00.—The A0620–00 region was observed for 45 ks

¹Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA.
 mgarcia@cfa.harvard.edu, jmcclintock@cfa.harvard.edu, rnarayan@cfa.harvard.edu, smurray@cfa.harvard.edu

²University College, Cork, Ireland, paulc@ucc.ie

TABLE 1
CHANDRA OBSERVATIONS OF QUIESCENT BHXN (DETECTED FLUXES AND LIMITS)

System	Instrument	Counts	Exposure ks	f_x ergs s ⁻¹ cm ⁻²	L_x ergs s ⁻¹	D kpc	m_1 $M_\odot$	log N_H cm ⁻²
GRO J0422+32	ACIS-I	16	18.8	6.6×10^{-15}	7.6×10^{30}	2.6	12	21.3
A0620-00 (XN Mon 75)	ACIS-S	123	38.3	1.8×10^{-14}	2.8×10^{30}	1.0	6.1	21.3
GS 2000+25 (XN Vul 88)	ACIS-I	5	21.5	2.4×10^{-15}	2.4×10^{30}	2.7	8.5	21.9
4U1543-47	ACIS-I	< 5	9.9	$< 4.2 \times 10^{-15}$	$< 3.0 \times 10^{31}$	6.1 ^a	6 ^a	21.6 ^b
GRO J1655-40	ACIS-S	65	42.5	1.2×10^{-14}	2.4×10^{31}	3.2	7	21.8
V404 Cyg (GS2023+338)	ACIS-S	1655	10.3	1.5×10^{-12}	4.9×10^{33}	3.5	12	22.0

Note: Values of N_H are taken from G98 and NGM; distances and primary masses are from M99, unless explicitly mentioned. Detected ACIS-S fluxes are over a 0.3–7.0 keV band; ACIS-I fluxes are over a 0.5–7.0 keV band; and emitted luminosities are over a 0.5–10.0 keV band.

^aOrosz et al 2001.

^b N_H from Greiner, Predehl & Harmon (1994) and van der Woerd, White & Kahn (1989).

on 2000 Feb 29 and 38.3 ks of low background data was obtained. A0620-00 was placed on the ACIS-S3 CCD (Garmire et al 1992) in order to maximize the counting rate from the possibly soft X-ray spectrum (Narayan, Barret, & McClintock 1997b). The CCD recorded 123 photons at a position consistent with the optical position of A0620-00.

GS 2000+25.— The GS 2000+25 region was observed with the Chandra ACIS-I detector on 1999 Nov 5 for a total of 21.5 ks. The CCD array recorded 5 photons near the position of GS 2000+25, which is a weak but significant detection. In order to more accurately determine the position of this source, we cross correlated the positions of all sources in the image with >10 counts against the USNO-A2 catalog. We found 4 matches within a 2'' search radius. We then used the IRAF ctio.coords task to determine the WCS for the X-ray image using these 4 stars. The plate solution has an RMS of 0.3'', and shows that the ICRS position of this weak source is RA=20:02:49.52, DEC=+25:14:10.34. Our ability to determine the centroid of this source is limited by the low number of counts to $\sim 0.5''$.

We determined the position of the optical counterpart of GS 2000+25 in a similar way, matching 8 USNO-A2 stars against a K-band image of the field we acquired at the LICK 3m with the Gemini camera. This plate solution has an RMS of 0.2'', and shows the position of the optical (IR) counterpart to be RA=20:02:49.55, DEC=+25:14:10.94. The offset between the two positions is $0.7'' \pm 0.8''$, thus the position of the weak X-ray source is consistent with GS 2000+25.

4U 1543-47.— A 9.9 ks ACIS-I observation of 4U 1543-47 took place on 2000 July 26, and failed to detect even a single photon from this source. We compute a conservative upper limit to the luminosity by assuming a that < 5 photons were detected. This luminosity (see Table 1) is a factor of ~ 100 below the previous upper limit (M99).

GRO J1655-40.— A 42.5 ks ACIS-S observation of GRO J1655-40 took place on 2000 July 2, and detected 65 photons from this source. We note that this corresponds to a luminosity a factor of ~ 10 below the previously measured quiescent luminosity (G98). Given the quiescent variations seen in V404 Cyg (Wagner et al 1994), this could simply be indicative of typical quiescent variations. Alternately, we note that the previous quiescent observations were in between two large outbursts sepa-

rated by ~ 1 year, and therefore may not have been indicative of the true quiescent level. The Chandra observations reported here occurred ~ 4 years after the last outburst, and therefore may accurately measure the quiescent level.

V404 Cyg.— A 10.3 ks ACIS-S observation of V404 Cyg took place on 26 April 2000, and allowed detection of 1655 photons from this source.

3. DISCUSSION

We have collected in Table 2 the available quiescent X-ray luminosities and upper limits for BHXN and NSXN. We have supplemented our Chandra measurements with the recent quiescent detections of SAX J1808.4-3658 (Stella et al 2000; Dotani, Asai & Wijnands 2000) and included previous data compiled by M99. Figure 1 displays the Eddington-scaled luminosities as a function of orbital period P_{orb} . For calculating L_{Edd} , we used the BH mass estimates in Table 1 and assumed that all NSs have a mass of $1.4M_\odot$.

With the Chandra detections of GRO J0422+32 and GS 2000+25 reported here, three short period BHXN have now been detected in quiescence. A greatly improved upper limit for a fourth short period system, 4U1543-47, is nearly comparable to these detections. In addition, we find a new, ten-times fainter quiescent luminosity for GRO J1655-40, which has a period intermediate between the NSXN and the long period BHXN V404 Cyg.

The new data points significantly strengthen our earlier claim (NGM, G98, M99) that BHXN have much lower (roughly a factor of 100) quiescent luminosities than NSXN. As explained in §1, such a difference in luminosity is natural if quiescent accretion proceeds via a radiatively inefficient ADAF and if the primaries in BHXN have event horizons. Thus, the new data bolster the evidence for event horizons in BHXN.

Our argument relies on the reasonable assumption that the Eddington scaled mass accretion rate is similar in quiescent BHXN and NSXN. At the short orbital periods characteristic of the NSXN in our sample, angular momentum loss through gravitational radiation is expected to be the dominant mechanism driving mass transfer from the secondary. The Eddington-scaled mass transfer rates are then likely to be roughly similar in BHXN and NSXN of similar P_{orb} (M99). At the long period

of V404 Cyg and perhaps GRO J1655–40, nuclear evolution is expected to drive the mass transfer rate to substantially higher values, and so these systems are less useful for our purposes. Another reason for ignoring these systems is the fact that there are no NSXN with similar P_{orb} for comparison.




Figure 1 (Top): Quiescent luminosities of BHXN (filled circles) and NSXN (open circles) from Table 2. Data points not included in M99 are circled. Multiple quiescent detections are included. (Bottom): Shows only the lowest quiescent detections or Chandra upper limits.

We also assume that f_* , the fraction of matter transferred from the secondary which actually reaches the central star, is the same in BHXN and NSXN. Out-flowing winds from an ADAF (Narayan & Yi 1994, 1995, Blandford and Begelman 1999, M99) tend to reduce f_* , but there is no reason to expect winds to be stronger (by a factor of 100) in BHXN than in NSXN. In fact, a centrifugal propeller and/or radio pulsar action could reduce f_* in NSXN without affecting BHXN (NGM, M99). Therefore, if at all, we expect f_* to be lower in NSXN than in BHXN, rather than the other way round. This strengthens the case for event horizons in BHXN.

In this connection, we note that SAX J1808.4-3658 is the least luminous of the NSXN. It also may have a time averaged mass transfer rate that is lower than that typical of XN,

perhaps due to the action of the radio pulsar on the secondary (Chakrabarty & Morgan 1998). Given that SAX J1808.4-3658 is the only example of a NSXN which shows permanent coherent pulsations, it seems possible that it may also have an unusually efficient “propeller.” In addition, this system may have already evolved past the minimum orbital period and the orbit may now be expanding (King 2000; Dotani et al 2000). Despite all these anomalies, this NSXN is ~ 60 times brighter than our three short period BHXN (see Figure 2).

Table 2
Quiescent Luminosities

System	P_{orb} (hr)	$\log[L_{min}]$ (erg s^{-1})
(1)	(2)	(3)
○ SAX J1808-365	2.0 ^a	32.0^b, 31.5^c
○ EXO 0748-676	3.82	34.1
○ 4U 2129+47	5.2	32.8
○ Cen X-4	15.1	32.4
○ Aql X-1	19	32.6
○ H1608-52	12^d	33.3
● GRO J0422+32	5.1	30.9, < 31.6
● A0620-00	7.8	30.5, 30.8^e
● GS 2000+25	8.3	30.4, < 32.2
● GS1124-683	10.4	< 32.4
● H1705-250	12.5	< 33.0
● 4U 1543-47	27.0	< 31.5, < 33.3
● GRO J1655-40	62.9	31.3, 32.4
● V404 Cyg	155.3	33.7, 33.1^f

NOTES — References are as in M99, unless otherwise noted. New and/or CHANDRA Measurements are in **BOLD FACE**. (1) ○ indicates a NS primary and ● a BH primary. (2) Orbital periods. (3) Luminosities in quiescence in the 0.5-10 keV band (corrected for the revised distances).

^aChakrabarty and Morgan 1998.

^bStella et al 2000; assuming $D = 2.5$ kpc.

^cDotani, Asai, and Wijnands 2000.

^dWachter 2000 reports a new 12 hr period, intermediate between the previously reported 98.4 hr (Ritter & Kolb 1998) and 5 hr (Chen et al. 1998) periods.

^eRecomputed for $\alpha = 2$ from NGM.

^fKong 2000.

Campana & Stella (2000) argue that it is not correct to compare only X-ray luminosities, as we have done, but that we should include also the quiescent non-stellar optical and UV luminosity. This point is not obvious since the origin of the optical/UV luminosity is presently unclear. Within the ADAF model, it depends on the poorly known radius at which the inner edge of the accretion disk evaporates into the ADAF (e.g., Narayan, McClintock & Yi (1996); Narayan, Barret & McClintock 1997b). It also depends on the strength of winds from the ADAF (Quataert & Narayan 1999). If the non-stellar optical/UV luminosity originates in the outer accretion disk, or in the hot spot where the mass transfer stream from the secondary impacts the disk, then the fact that the optical/UV luminosities of BHXN and NSXN are similar (Campana & Stella 2000) provides observational confirmation that the mass transfer rates in the two kinds of system are similar, as we have assumed. An

0.25 mag modulation in the far-UV flux of A0620–00 on an orbital time scale suggests that the far-UV flux may be modulated with the orbital phase (McClintock 2000), which indicates an origin in the hot spot.

Bildsten & Rutledge (2000, but see Lasota 2000) suggest that the X-rays in BHXN may be produced by a rotationally enhanced stellar corona in the secondary, especially in the short period BHXN with the lowest luminosities. The detection of GRO J0422+32 with a luminosity of nearly 10^{31} ergs $^{-1}$ rules out the coronal interpretation for this system (Lasota 2000), and perhaps all the other BHXN in our sample as well. Also, the Chandra spectra of A0620–00 and GRO J1655–40 do not appear to be consistent with the stellar corona interpretation (McClintock et al 2001). If some (or all) of the X-ray emission in BHXN were coronal, then the accretion luminosities of the black holes would be even lower than our estimates and the argument for event horizons would be further strengthened.

A critical element in our comparison of NSXN and BHXN quiescent luminosities is the assumption that the quiescent X-ray luminosities of NSXN result from accretion. Brown, Bildsten & Rutledge (1998) have suggested that the luminosity could be due to crustal heating of the NS during outburst followed by cooling in quiescence. The rapid variability of the prototypical NSXN Cen X-4 (Campana et al 1997) is hard to explain in a cooling model, and shows that at most $\sim 1/3$ of the quiescent luminosity of this source is due to crustal cooling (M99). In addition, Cen X-4 and Aql X-1 have substantial power-law tails in their spectra (carrying about half the total luminosity), and it is hard to explain this spectral component with a cooling model. It thus seems reasonable to assume that accretion accounts for a substantial fraction of the quiescent X-

ray luminosity in most NSXN. The optical variability of NSXN in quiescence (McClintock & Remillard 2000; Jain et al 2000; Illovaisky & Chevalier 2000) provides ample evidence that accretion continues during quiescence.

Regardless of the caveats mentioned above, ultimately the dramatic difference in quiescent X-ray luminosities of BHXN and NSXN needs to be explained. Any explanation is likely to require a fundamental difference in the nature of the primaries in BHXN and NSXN. In our view, any straightforward explanation, whether based on ADAFs or not, will require postulating an event horizon in BHXN.

The Eddington scaled luminosities of some supermassive black holes (SMBH) in galactic nuclei (e.g., Garcia et al 2000; Baganoff et al 2000) are much smaller than those of the “stellar mass” black holes discussed here. Could this be evidence for the event horizon (Narayan et al. 1998)? Unfortunately, the mass accretion rates of SMBH are difficult to determine observationally (DiMatteo 2000; DiMatteo, Carilli, & Fabian 2000; Quataert, Narayan, and Reid 1999), and one cannot exclude the possibility that the low luminosities of SMBH are simply the result of extremely low accretion rates. In the case of quiescent BHXN, we have the luxury of a control sample of NSXN. By comparing quiescent BHXN and NSXN with similar orbital periods, we eliminate the uncertainty in the mass accretion rate, and thus obtain more secure evidence for the event horizon.

This work was supported in part by NASA Contract NAS8-39073 to the Chandra X-ray Center, contract NAS8-38248 to the HRC Team, and NSF grant AST 9820686 to RN. We thank A. Kong for helpful comments.

REFERENCES

- Arnaud, K., and Dorman, B., 2000, “XSPEC, An X-ray Spectral Fitting Package, Users Guide for version 11”, NASA/GSFC, available at: <http://heasarc.gsfc.nasa.gov/docs/xanadu/xspec/index.html>
- Baganoff, F. et al 2000, Poster paper at 2000 HEAD meeting, American Astronomical Society, HEAD meeting 32, 02.10
- Barret, D., McClintock, J.E., and Grindlay, J., 1996 ApJ 473, 963
- Bildsten, L., and Rutledge, R.E., 2000 ApJ 541, 908
- Brown, E.F., Bildsten, L. & Rutledge, R.E., 1998, ApJ Lett., 504, L95.
- Blandford, R.D. & Begelman, M.C., 1999 MNRAS 303L, 1B
- Campana, S., Mereghetti, S., Stella, L. & Colpi, M., 1997, A&A, 32 4, 941.
- Campana, S. 1998 ApJ 499, L65
- Campana, S., Stella, L., 2000 ApJ 541, 849
- Chakrabarty, D.; Morgan, E. H. 1998 Nature 394, 346
- Chen, W. et al., 1998, in *Accretion Processes in Astrophysics - Some Like it Hot*, eds. S. Holt & T. Kallman (Woodbury, NY: AIP), p. 347.
- Chandra X-ray Center (CXC) 2000a, “The CXC Data Manipulation User’s Guide”, Version 1.1, available at: http://asc.harvard.edu/ciao/download/doc/manual_dm.ps
- CXC, Chandra MSFC, and Chandra IPI Teams 2000b, “Chandra Proposers Observatory Guide (POG)” Version 2.0, available at: <http://asc.harvard.edu/udocs/docs/docs.html>
- DiMatteo, T., Carilli, C.L., & Fabian, A.C. 2000 astro-ph/0005516
- DiMatteo, T. 2000, submitted to ApJ
- Dotani, T., Asai, K., Wijnands, R., 2000 ApJ 543, L145
- Filippenko, A.V., Matheson, T., Ho, L.C. 1995 ApJ 455, 614
- Garcia, M.R., McClintock, J.E., Narayan, R., & Callanan, J., 1998, in the *Proceedings of the 13th NAW on CVs, Jackson Hole, WY*, eds. S. Howell, E. Kuulkers, and C. Woodward, p. 506 (G98)
- Garcia, M.R., Murray, S.S., Primini, F.A., Forman, W., Jones, C. 2000, ApJ 537 L23.
- Garmire, G.P. Ricker, G., Bautz, B., Burrows, D., Collins, S., Doty, J., Gendreau, K., Lumb, D., and Nousek, J. 1992 in “American Institute of Aeronautics and Astronautics Conference”, 8
- Greiner, J. Predehl, P. & Harmon, B.A. 1994, in *AIP Conf. Proc. 304, 2d Compton Symp.*, ed. C.E. Fichtel, N. Gehrels, & J.P. Norris (New York: AIP), 314
- Illovaisky, S.A. & Chevalier, C. 2000 IAUC 6025
- Jain, R. et al 2000, IAUC 7429
- King, A.R. 2000 MNRAS 315, L33
- Kong, A., 2000, PhD Thesis, Oxford University, UK.
- Lasota, J.P., 2000 A&A 360, 575
- McClintock, J.E., Garcia, M.R., Narayan, R., Murray, S.S. 2001, in prep
- McClintock, J.E. 2000, talk at Rossi 2000 Symposium at GSFC, March 22-24, 2000
- McClintock, J.E., & Remillard, R.R. 2000 ApJ 531, 956
- McClintock, J.E., Horne, K., and Remillard, R.A. 1995 ApJ 442, 358
- Menou, K., Esin, A.A., Narayan, R., Garcia, M.R., Lasota, J.P., & McClintock, J.E. 1999 ApJ 520, 276 (M99)
- Narayan, R. and Yi, I., 1994 ApJ 428, L13
- Narayan, R. and Yi, I., 1995 ApJ 444, 231
- Narayan, R., McClintock, J.E., & Yi, I. 1996 ApJ 457, 821
- Narayan, R., Garcia, M.R., & McClintock, J.E. 1997a ApJ 478, L79 (NGM)
- Narayan, R., Barret, D., & McClintock, J.E. 1997b ApJ 482, 448.
- Orosz, J. et al 2001, in prep
- Plucinsky, P.P. & Virani, S.N. 2000 in *SPIE X-Ray Optics, Instruments, and Missions*, ed J. Trümper & B. Aschenbach, Proc. SPIE 4012, 669
- Predehl, P. & Schmitt, J.H.M.M. 1995 A&A 293, 889.
- Quataert, E., Narayan, R., & Reid, M.J. 1999 ApJ 517, L101
- Quataert, E., and Narayan, R., 1999 ApJ 520, 298
- Stella, L., Campana, S., Mereghetti, S., Ricci, D. and Israel, G.L. 2000 ApJ 537, L115
- Ritter, H. & Kolb, U., 1998, A&AS, 129, 83
- van der Woerd, H., White, N.E., & Kahn, S.M. 1989, APJ, 344, 320
- Van Speybroeck, L., Jerius, D., Edgar, R.J., Gaetz, T.J., Zhao, P., and Reid, P.B., 1997 Proc SPIE 3113, 98
- Vikhlinin, A., McNamara, B. R., Forman, W., Jones, C., Quintana, H., Hornstrup, A. 1998 ApJ 502, 558.
- Wagner, M.R., Starrfield, S.G., Hjellming, R.M., Howell, S.B., Kreidl, T.J., 1994 ApJ 429 L25
- Wachter, S., 2000, in “1999 Pacific Rim Conference on Stellar Astrophysics”, ed K.S. Cheng, (Kluwer:Dordrecht) p. 451
- Weisskopf, M.C., and O’Doell, S.L. 1997 Proc. SPIE 3113, 2